


## Sideritis cystosiphon

### 1. National level

Biogeographical regions and/or marine regions concerned within the Member State: **MAC**

map-range

map-distribution



map-favourable-range



### 2. Biogeographical or marine level

2.1 Biogeographical region or marine region: **MACARONESIAN**

2.2 Published sources and/or websites:

Beltrán E., Wildpret W., León C., García A. & A. Reyes, 1999. Libro Rojo de la Flora Canaria contenida en la Directiva-Hábitats Europea. Ministerio de Medio Ambiente. 694 pp.

Bañares A., Blanca G., Güemes J., Moreno J.C. & Ortiz S., eds. (2003). Atlas y Libro Rojo de la Flora Vascular Amenazada de España. Dirección General de Conservación de la Naturaleza. Madrid. 1072 pp

[http://www.mma.es/porta/cciones/biodiversidad/inventarios/inb/flora\\_vascular/pdf/736.pdf](http://www.mma.es/porta/cciones/biodiversidad/inventarios/inb/flora_vascular/pdf/736.pdf)

2.3 Range of the species type in the biogeographic region or marine region

- |   |                |
|---|----------------|
| 2.3.1 Surface area of species range in km2: | 100            |
| 2.3.2 Date of range determination:          | 2003           |
| 2.3.3 Quality of data concerning range:     |                |
| 2.3.4 Range trend:                          | Increasing (+) |

## Sideritis cystosiphon

2.3.5 Range trend magnitude in km2 (optional):

2.3.6 Range trend period: 1990-2003

2.3.7 Reasons for reported trend: Improved knowledge/more accurate data  
and/or specify

### 2.4 Population of the species in the biogeographic region or marine region

2.4.1 Population size estimation:

Population size estimation (minimum)	Population size estimation (maximum)	Population units
884	884	Number of individuals

2.4.2 Date of population estimation: 2003

2.4.3 Methods used for population estimation: From comprehensive inventory

2.4.4 Quality of data on area: Good e.g based on extensive surveys

2.4.5 Population trend: Increasing (+)

2.4.6 Population trend magnitude (km2):

2.4.7 Population trend period: 1990-2003

2.4.8 Reasons for reported trend: Improved knowledge/more accurate data  
and/or specify:

2.4.9 Justification of % thresholds for trends (optional):

2.4.10 Main pressures: 502 - roads, motorways  
790 Other pollution or human impacts/activities  
965 - predation

2.4.11 Threats 502 - roads, motorways  
965 - predation

### 2,5 Habitat for the species in the biogeographic region or marine region

2.5.1 Habitats for the species: Especie xerófila y heliófila que crece en el dominio de los cardonales del piso bio

2.5.2 Area estimation (km2): 25

2.5.3 Date of estimation: 2007

2.5.4 Quality of the data: Moderate e.g. based on partial data with some extrapolation

2.5.5 Trend of the habitat: Unknown (X)

2.5.6 Trend period: 1970-2006

2.5.7 Reasons for reported trend:

Other (specify):

2.6 Future prospects for the species: Good prospects - species expected to survive and prosper

### 2.7 Complementary information

2.7.1 Favourable reference range (km2): 25

2.7.2 Favourable reference population: 25

2.7.3 Suitable habitat for the species (km2):

2.7.4 Other relevant information (optional): Favourable reference area: 200 individuals/population Una carretera pasa por la

Conclusion	Biogeographical or marine level	Conclusions within Natura 2000 sites (optional)
Conclusions: (2.3) Range:	Inadequate but improving (U1+)	
Conclusions: (2.4) Population:	Inadequate but improving (U1+)	

**Sideritis cystosiphon**

Conclusions: (2.5) Habitat for the species:	Unknown (XX)
Conclusions: (2.6) Future prospects:	Inadequate but improving (U1+)
Conclusions: Overall assessment:	Inadequate but improving (U1+)