


## Teline salsoloides

### 1. National level


Biogeographical regions and/or marine regions concerned within the Member State: **MAC**

map-range

map-distribution



map-favourable-range



### 2. Biogeographical or marine level

2.1 Biogeographical region or marine region: **MACARONESIAN**

2.2 Published sources and/or websites:

Beltrán E., Wildpret W., León C., García A. & A. Reyes (1999). Libro Rojo de la Flora Canaria contenida en la Directiva-Hábitats Europea. Ministerio de Medio Ambiente. 694 pp.

Bañares A., Blanca G., Güemes J., Moreno J.C. & Ortiz S., eds. (2003). Atlas y Libro Rojo de la Flora Vascular Amenazada de España. Dirección General de Conservación de la Naturaleza. Madrid. 1072 pp

[http://www.mma.es/porta1/secciones/biodiversidad/inventarios/inb/flora\\_vascular/pdf/853.pdf](http://www.mma.es/porta1/secciones/biodiversidad/inventarios/inb/flora_vascular/pdf/853.pdf)

2.3 Range of the species type in the biogeographic region or marine region

2.3.1 Surface area of species range in km2: 50

2.3.2 Date of range determination: 2003

2.3.3 Quality of data concerning range:

2.3.4 Range trend: Stable (=)

# Teline salsoloides

- 2.3.5 Range trend magnitude in km2 (optional):
- 2.3.6 Range trend period: 1993-2003
- 2.3.7 Reasons for reported trend: Natural processes
- and/or specify

## 2.4 Population of the species in the biogeographic region or marine region

- 2.4.1 Population size estimation:
- Population size estimation (minimum)

Population size estimation (maximum)

Population units

156

156

Number of individuals
- 2.4.2 Date of population estimation: 2003
- 2.4.3 Methods used for population estimation: From comprehensive inventory
- 2.4.4 Quality of data on area: Good e.g based on extensive surveys
- 2.4.5 Population trend: Increasing (+)
- 2.4.6 Population trend magnitude (km2):
- 2.4.7 Population trend period: 1993-2003
- 2.4.8 Reasons for reported trend: Improved knowledge/more accurate data
- and/or specify:
- 2.4.9 Justification of % thresholds for trends (optional):
- 2.4.10 Main pressures: 965 - predation
- 2.4.11 Threats 965 - predation

## 2,5 Habitat for the species in the biogeographic region or marine region

- 2.5.1 Habitats for the species: Cantiles litorales de elevada pendiente, escaso suelo y acusada inestabilidad geol
- 2.5.2 Area estimation (km2): 50
- 2.5.3 Date of estimation: 2007
- 2.5.4 Quality of the data: Moderate e.g. based on partial data with some extrapolation
- 2.5.5 Trend of the habitat: Stable (=)
- 2.5.6 Trend period: 1970-2006
- 2.5.7 Reasons for reported trend: NaturalProcesses
- Other (specify):

- 2.6 Future prospects for the species: Poor prospects - species likely to struggle unless conditions change

## 2.7 Complementary information

- 2.7.1 Favourable reference range (km2): 25
- 2.7.2 Favourable reference population:
- 2.7.3 Suitable habitat for the species (km2): 25
- 2.7.4 Other relevant information (optional): Favorable reference population: 150 individuals/population. La mayor parte de la

Conclusion	Biogeographical or marine level	Conclusions within Natura 2000 sites (optional)
Conclusions: (2.3) Range:	Bad (U2)	
Conclusions: (2.4) Population:	Bad (U2)	
Conclusions: (2.5) Habitat for the species:	Inadequate (U1)	
Conclusions: (2.6) Future prospects:	Bad (U2)	
Conclusions: Overall assessment:	Bad (U2)	