


Dendriopoterium pulidoi

1. National level

Biogeographical regions and/or marine regions concerned within the Member State: **MAC**

map-range

map-distribution



map-favourable-range



2. Biogeographical or marine level

2.1 Biogeographical region or marine region: **MACARONESIAN**

2.2 Published sources and/or websites:

Beltrán E., Wildpret W., León C., García A. & A. Reyes (1999). Libro Rojo de la Flora Canaria contenida en la Directiva-Hábitats Europea. Ministerio de Medio Ambiente. 694 pp.

Bañares A., Blanca G., Güemes J., Moreno J.C. & Ortiz S., eds. (2003). Atlas y Libro Rojo de la Flora Vascular Amenazada de España. Dirección General de Conservación de la Naturaleza. Madrid. 1072 pp

http://www.mma.es/porta1/secciones/biodiversidad/inventarios/inb/flora_vascular/pdf/1104.pdf

2.3 Range of the species type in the biogeographic region or marine region

2.3.1 Surface area of species range in km2: 200

2.3.2 Date of range determination: 2003

2.3.3 Quality of data concerning range:

2.3.4 Range trend: Increasing (+)

Dendriopoterium pulidoi

2.3.5 Range trend magnitude in km2 (optional):

2.3.6 Range trend period: 1995-2003

2.3.7 Reasons for reported trend: Improved knowledge/more accurate data
and/or specify

2.4 Population of the species in the biogeographic region or marine region

2.4.1 Population size estimation:

Population size estimation (minimum)	Population size estimation (maximum)	Population units
5700		Number of individuals

2.4.2 Date of population estimation: 2003

2.4.3 Methods used for population estimation: From comprehensive inventory

2.4.4 Quality of data on area: Moderate e.g. based on partial data with some extrapolation

2.4.5 Population trend: Increasing (+)

2.4.6 Population trend magnitude (km2):

2.4.7 Population trend period: 1995-2003

2.4.8 Reasons for reported trend: Improved knowledge/more accurate data
and/or specify:

2.4.9 Justification of % thresholds for trends (optional):

2.4.10 Main pressures: 965 - predation

2.4.11 Threats 965 - predation

2,5 Habitat for the species in the biogeographic region or marine region

2.5.1 Habitats for the species: Crece en paredones de tobas ignimbríticas y coladas riolítico-traquíticas peralcali

2.5.2 Area estimation (km2): 225

2.5.3 Date of estimation: 2007

2.5.4 Quality of the data: Moderate e.g. based on partial data with some extrapolation

2.5.5 Trend of the habitat: Stable (=)

2.5.6 Trend period: 1970-2006

2.5.7 Reasons for reported trend: NaturalProcesses

Other (specify):

2.6 Future prospects for the species: Good prospects - species expected to survive and prosper

2.7 Complementary information

2.7.1 Favourable reference range (km2): Less than

2.7.2 Favourable reference population: Less than

2.7.3 Suitable habitat for the species (km2): 125

2.7.4 Other relevant information (optional): En general, las poblaciones muestran cohortes de diferentes edades indicando cie

Conclusion	Biogeographical or marine level	Conclusions within Natura 2000 sites (optional)
Conclusions: (2.3) Range:	Inadequate but improving (U1+)	
Conclusions: (2.4) Population:	Inadequate but improving (U1+)	
Conclusions: (2.5) Habitat for the species:	Inadequate but improving (U1+)	
Conclusions: (2.6) Future prospects:	Inadequate but improving (U1+)	
Conclusions: Overall assessment:	Inadequate but improving (U1+)	